

Econometrics and the Science of Climate Change

19th May 2011

Timothy Curtin¹

1. Introduction

Econometrics has a long history as the technique of choice for testing the merits of alternative hypotheses across most of the social sciences as well as many of the natural and materials sciences, not to mention pharmaceutical science, where it is widely used to evaluate the efficacy of alternative medications, including the use of placebos as counterfactuals. It is acknowledged by most climate scientists that the atmospheric concentration of carbon dioxide (hereafter denoted $[\text{CO}_2]$) and other greenhouse gases (in aggregate denoted as $[\text{CO}_{2e}]$) is not necessarily the only determinant of climate change, as they accept that there are natural factors also at work. Nevertheless one will in vain search the literature of climate science for any evidence of the use of econometrics to test the core hypothesis that “most” of the temperature change observed over the last century is attributable to the build-up in the atmosphere of anthropogenic emissions of CO_{2e} , of which CO_2 is by far the largest in volume terms. In particular none of the leading texts such as the IPCC’s Solomon *et al.* (2007), Stern (2006) and Garnaut (2008, 2011) performs or reports any econometric analysis of their core hypothesis. This chapter seeks to begin filling that gap, and finds that the core hypothesis is falsified at a wide variety of locations with lengthy time series data on various climatic variables, including atmospheric water vapour (i.e. $[\text{H}_2\text{O}]$), opacity of the sky (OPQ), and solar radiation (SR) received at the earth’s surface (as opposed to the top of the atmosphere). Multi-variate econometric analysis shows that at none of these locations is the role of $[\text{CO}_2]$ statistically significant, and even that it is can be negatively correlated with changes in temperature, whereas these other variables play highly significant roles. If the core hypothesis of climate science cannot be confirmed at any specific location, then by Popper’s Black Swan paradigm, it cannot be confirmed for the globe, as that is the average of the local.

2. Air, Water, and Climate Change

In 1853-54 there was a serious outbreak of cholera in London’s Soho district. The prevailing view of scientists at the time was that cholera - like climate change now - was caused by an invisible miasma in the air, and those espousing the conventional wisdom of the time all deferred to various Royal Societies (e.g. the Royal College of Physicians) as the definitive authorities. John Snow, a doctor who had pioneered the use of chloroform, in 1849 published at his own expense the first book *On the mode of communication of cholera* to challenge the conventional wisdom of the authorities of the day, and was duly put down by them. Had Snow derived only the average number of cholera-related deaths in each street in Central London he would not have been able to challenge the orthodox view. Snow instead

¹ Timothy Curtin is a retired former academic economist (1964-1970) and economic adviser (1970-2002). He lives in Canberra, Australia. Email: tcurtin@bigblue.net.au.

showed how the incidence of cholera in the 1854 outbreak was closely correlated with the nature of the drinking water supplied by the Southwark and Vauxhall water company on the one hand, and the Lambeth company on the other, to individual houses on the same streets. He eventually won acceptance for his contention that cholera is a water-borne disease, having shown *statistically* how most cholera deaths occurred in premises taking their water from the first company, which had its water intakes adjacent to sewerage outlets to the Thames, while very few deaths occurred in premises whose water was supplied by the Lambeth company, which in 1849 had moved its intake to upstream from the sewerage outlets.³

It is fair to say that the majority of climate scientists mostly content themselves by comparison of global mean temperature trends with the trend in the atmospheric concentration of carbon dioxide and other greenhouse gases, with some excursions into average temperatures of land and sea surfaces or of the northern and southern hemispheres. Solomon *et al.* (2007) summarise these aggregative exercises. This chapter follows the Snow practice, of examining not streets but houses within each street, by examining a wide range of climatic variables at specific locations in the USA, from Barrow in the Arctic to Hawaii near the equator to New York in the north-east. Like Snow, it finds that in every case water vapour is a far more significant determinant of temperature change than atmospheric carbon dioxide.

Snow's pioneering counterfactual analysis has recently been cited in two advanced econometrics textbooks (Angrist and Pischke 2009, Freedman 2010) that explain how multivariate regression analysis can be used to evaluate competing theories of causation. This chapter shows how such analysis reveals that changes in atmospheric water vapour and *in situ* solar radiation (SR) are much more powerful explanations of climate change than changes in carbon dioxide levels, because like Snow it uses counterfactuals to show that although the atmospheric concentration of carbon dioxide is ubiquitous, temperature changes are not, and that in most places changes in atmospheric water vapour and SR have a very much larger – and much more statistically significant – association with changes in temperature. Were he alive now it seem more than likely John Snow would have been as sceptical of airborne carbon dioxide being the cause of changing climate everywhere as he was of miasmatic air being guilty of spreading cholera.

The Australian Government's chief climate change adviser has since 2008 been the economist Ross Garnaut. His first report to the Government (2008) strongly argued that the government needed to introduce a "carbon pollution reduction scheme". Its attempt to do so was defeated in the country's Parliament in late 2009, and that led in part to the ousting of the prime minister, Kevin Rudd, in mid-2010. The new prime minister, Julia Gillard

³ Laveran and Ross, discoverers of the parasitic nature of malaria, had similar experiences to Snow's. The weight of scientific opinion at the French and British academies and elsewhere was that like cholera, malaria (as its name implied) was an airborne miasmatic disease. Laveran and Ross suspected water-based mosquitos carried a parasite they passed on to those they bit, and like Snow they did counterfactual analysis until they found the particular species of mosquito that carried the parasite (Rocco 2003:250-280).

announced after the general election (October 2010) that a Carbon Tax would be introduced on 1st July 2012, and that the government would again be advised by Ross Garnaut. In early 2011 he produced 8 reports updating his 2008 Review, including his *The Science of Climate Change* (2011, hereafter SCC), which is notable for its avoidance of econometrics and counterfactuals, and concludes (in its section 4.2.2) by invoking the authority amongst others of the Royal Society (2010) and the Australian Academy of Science (Allison *et al.* 2010) with their support for the findings of the International Panel on Climate Change (IPCC) that “most” global warming is caused by airborne carbon dioxide (and some lesser so-called greenhouse gases).⁴

The IPCC’s *Climate Change 2007 The Physical Science Basis* goes out of its way (Solomon *et al.* 2007:28) to dismiss any role in climate change for the emissions of water vapour that are *simultaneous* with emissions of carbon dioxide whenever there is combustion of hydrocarbon fuels. Yet basic chemistry shows that while water vapour emissions are only in the range of 30-50 per cent of CO₂ emissions by weight, *in addition* to any moisture content of the fuel and to the steam produced during the power generation process, their effect on surface temperature is as much as twice as large pro rata.⁵ It follows that the Australian Government’s proposed tax on carbon, based on the *Garnaut Reviews* (2008, 2011) will also be a tax on increases in water vapour and thereby, unavoidably, on rainfall. The political implications of that remain to be played out, but it is characteristic of all the papers on climate change cited by the *Reviews* that they dwell only on the supposed – and far-off – external costs of hydrocarbon combustion and never mention the demonstrably larger immediate benefits of elevated atmospheric carbon dioxide, warmer temperatures, and increasing rainfall on the world’s primary production.

3. Statistics and the Science of Climate Change

Ross Garnaut’s SCC like his 2008 *Review* is notable as much for what it leaves out as for what it includes. For example, here is its first “key point” (2010:2):

Observations and research outcomes since 2008 have confirmed and strengthened the position that the mainstream science then held with a high level of certainty, that the Earth is warming and that human emissions of greenhouse gases are the primary cause...The statistically significant warming trend has been confirmed by observations over recent years: global temperatures continue to rise around the midpoints of the range of the projections of the Intergovernmental Panel on Climate Change (IPCC) and the presence of a warming trend has been confirmed.

⁴ Garnaut’s SCC implies that the Australian Academy of Science (AAS) report (Allison *et al.* 2010) provides independent support for the IPCC’s AR4 (2007), when most of its authors were themselves leading contributors to AR4 (Solomon *et al.* 2007).

⁵ Water-vapour feedback has “the capacity to about double the direct warming from greenhouse gas increases”, Dessler *et al.* (2008:1).

However, the paper *Global Temperature Trends* by Breusch and Vahid (2011) commissioned by Garnaut for his 2008 *Review*, and updated for SCC, does not claim to provide any support for a warming trend being attributable to human emissions of carbon dioxide.

We conclude that there is sufficient statistical evidence in the temperature data of the past 130-160 years to conclude that global average temperatures have been on a warming trend. The evidence of a warming trend is present in all three [sic] of the temperature series. Although we have used unit roots and linear trends as a coordinate system to approximate the high persistence and the drift in the data in order to answer the questions, *we do not claim that we have uncovered the nature of the trend in the temperature data*. There are many mechanisms that can generate trends and *linear trends are only a first order approximation...* It is impossible to uncover detailed trend patterns from such temperature records *without corroborating data* from other sources and close knowledge of the underlying climate system (2011:13, my emphasis).


Thus Breusch and Vahid explicitly exclude any assessment of the possible role of rising [CO₂] in whatever very slight global warming trend is evident in the observed data. Their finding of a rising temperature trend depends partly on the absence of the tropics from all global temperature sets for the period between 1850 and 1910, as it was not until the 1950s that global temperature becomes a valid statistic, for only then did global surface instrumental temperature coverage reach 80 per cent, and it is now well below that level again. Regrettably Breusch and Vahid do not assess the trends in the more truly global satellite temperature data sets, which surely should “corroborate” their surface trends data even though they do not go further back than 1978. The truth is shown in my Fig.1: the linear trend in the UAH satellites’ global data from December 1978 to February 2011 has an R² of 0.345, which although that is below the minimum for a good fit indicates a rise of 0.0012°C per month since 1978, or 0.0144 p.a., 0.144 per decade, and 1.44°C per century, well below the 3°C predicted by the IPCC, let alone the 5°C predicted for 2100 by Garnaut (2008:Fig.4.5) if there is no “mitigation” (i.e. reduction of projected Business as Usual emissions, BAU). The UAH data do *not* in fact “corroborate” the three surface temperature data sets used by Breusch and Vahid, for the period when they overlap.⁶ For example, their Gistemp data set shows a very much larger and apparently better fit linear trend since 1978 (see Fig.2), with the annual change in the Gistemp monthly data (1979-2010) at 0.0196°C, which is 36 per cent higher than the UAH trend of 0.0144 °C.

However, the AAS (Allison *et al.* 2010) asserts there is no disagreement between satellite and surface temperature records - “while a disagreement did exist in the 1990s, it has largely been resolved by correction of biases in the satellite data, for example to account for drift in

⁶ “The raw surface temperature data from which all of the different global surface temperature trend analyses are derived are essentially the same. The best estimate that has been reported is that 90–95% of the raw data in each of the analyses is the same (P. Jones, personal communication, 2003)” (Pielke snr. *et al.* 2007:13). See Stockwell (2009) for a comment on Rahmstorf *et al.* 2007 and the pitfalls of using the surface sets to determine apparent breaks in trends.

satellite orbits over time. Given the remaining uncertainties in satellite-derived trends, there is now acceptable agreement between satellite and ground-based measurements of surface temperature (Box 5). However Klotzbach *et al.* confirm my finding that “Surface temperature data sets from the National Climate Data Center and the Hadley Center show larger trends over the 30-year period than the lower-tropospheric data from the University of Alabama in Huntsville and Remote Sensing Systems data sets. The differences between trends observed in the surface and lower-tropospheric satellite data sets are statistically significant in most comparisons, with much greater differences over land areas than over ocean areas. These findings strongly suggest that there remain important inconsistencies between surface and satellite records” (2009:Abstract).


Fig. 1 Global monthly mean temperature UAH satellite data



Source: <http://www.drroyspencer.com/2011/01/dec-2010-uah-global-temperature-update-0-18-deg-c/>

Note: Linear Global Monthly Mean Temperature Trends are 0.0144°C p.a., 0.144°C per decade, and 1.44°C per 100 years (the anomaly is departure from 1981 to 2010 average in °C).

Fig.2 Annual Surface Instrumental Global Mean Temperatures Anomalies



Source: <http://data.giss.nasa.gov/gistemp>

Note: The Gistemp anomalies are expressed in 0.01°C divergences from the global mean for 1951-1980, so need to be divided by 100 to get °C. The linear trends are 0.019°C p.a, 0.196°C per decade, and 1.96 °C to 2110.

Garnaut’s *Update Reports* (2011) and *2008 Review* are both in the nature of a Prospectus, inviting the Government and people of Australia to invest in a Carbon Tax (and eventual Emissions Trading Scheme, ETS) that will save them the costs of “dangerous climate change”, but “a prospectus is required to contain **all the information** that investors and their advisers would require and expect to make an informed assessment of the offer being made”, *Australian Stock Exchange 2011*, my emphasis). The absence of any mention let alone analysis in Breusch and Vahid (2008 and 2011) of the UAH and RSS satellite data sets in addition to the Gistemp/NCDS/HadleyCRU surface data sets is regrettable.

However, there is a further omission of “all the information investors and their advisers would require and expect” in both Garnaut *Reviews* when they make this statement:


No-mitigation case – based on no action undertaken to mitigate climate change, and used as a ‘reference’ to assess the benefits of climate change action that accrue from the climate change impacts that are avoided. By the end of the century the concentration of long-lived greenhouse gases in the atmosphere is **1565** ppm carbon dioxide equivalent (Garnaut 2011:5 and Fig.1, my emphasis).

The present (2011) atmospheric concentration of CO₂ is 390 parts per million (ppm) of the atmosphere. To this the *Reviews* add the atmospheric levels of other greenhouse gases, chiefly methane (CH₄) and nitrous oxide (N₂O), which are expressed in CO₂ “equivalent” amounts that bring the CO_{2e} concentration to between 455 and 465 ppm in 2010. To get from 460 ppm to 1565 ppm by 2100 requires that the rate of growth of CO_{2e} from now until 2100 has to be 1.0137 per cent p.a. The actual growth of atmospheric CO₂ from 1958 to 2010 was 0.295 per cent p.a., and projecting CO_{2e} at that rate (the growth rates of CH₄ and N₂O are much lower than that of CO₂, see IPCC, Solomon *et al.* 2007:141)⁸ to 2100 produces only 600 ppm of CO_{2e} by 2100, less than 38 per cent of the *Reviews*’ projection of 1565 ppm. The *Reviews* offer no basis for predicting any *acceleration* in the “no mitigation” BAU rate of growth of the atmospheric concentration of CO_{2e} above the actual rate from 1958 to 2010, other than stressing the undoubted increasing growth of manmade CO₂ emissions – and ignore the finding in Knorr (2009) that “despite the predictions of coupled climate-carbon cycle models, no trend in the airborne fraction [of CO₂ emissions] can be found”. Nor does the SCC mention that Knorr specifically rebuts Raupach *et al.* 2008, one of 7 almost identical papers by those authors that are cited by SCC for its claim that the ability of the worlds’ biospheres to continue absorbing around 44 per cent of CO₂ emissions is “declining” (see also Curtin 2009 for analysis of the raw data from 1958 to 2008 rebutting this claim).

4. The Science of Climate Change


Real science involves very precise formulae, like the formula for complete combustion of a typical hydrocarbon fuel (e.g. C₃H₈) in the presence of air (using pure oxygen tends to raise the proportion of H₂O in the outputs):

⁸ Growth of the methane concentration from 1998 to 2005 was NIL, despite SCC’s claim that its growth rate is increasing, and of N₂O, 11 per cent over that period (in total, not p.a.), against 13 per cent in CO₂ (IPCC, Solomon *et al.* 2007:141). “The major (95%) sink for methane is OH radicals in the troposphere. These are concentrated in the tropics as water vapour is their source and water vapour is at a maximum concentration in the Tropics. If there is a change in humidity at about the year 2000 then there should be a change in the atmospheric concentration of methane. An increase in humidity implies an increase in OH radicals and a consequent reduction of methane... For the period 1991 to 2009 the annual increase in methane falls from 5.8 +/- 0.4 ppb per year before 1999 to 1.3 +/- 0.4 ppb per year after 1999”, Quirk 2011, see also Chambers, 2010.



Climate science has never produced any such formulae, and formula (1) receives no mention in the IPCC's AR4 WG1 (Solomon *et al.*, 2007) despite its demonising of hydrocarbon combustion.

It is not surprising that just as climate science never divulges equation (1) despite it showing how CO₂ enters the atmosphere when there is combustion of hydrocarbons, it also never mentions (2), the formula whereby a large proportion of the emissions of CO₂ is absorbed by photosynthesis both on land and in the oceans:



Or, in words, carbon dioxide + water + light energy → carbohydrate + oxygen.¹⁰

It would seem that the presence of CO₂ and H₂O on the RHS of equation (1) and on the LHS of equation (2) are inconvenient truths for all climate scientists named in the IPCC's Solomon *et al.* (2007). Although there are three references to photosynthesis in that volume, none of them mention that its main output is the carbohydrates on the RHS of (2) which of course constitute the world's food supply (see below).

Climate science does not offer formulae like (1) and (2). The closest it gets to producing a formula is for the "radiative forcing" (RF) of greenhouse gases, namely the strength of their ability "forcibly" to radiate heat back to earth, are their equations (3) and (4). In (3) the change in equilibrium surface temperature (ΔT_s) arising from RF is given by:¹¹

$$\Delta T_s = \lambda \Delta F \dots(3)$$

where λ is the *assumed* (not proven) climate sensitivity, with units in K/(W/m²), and ΔF is the *assumed* radiative forcing, given by

$$\Delta F = 5.35 \times \ln (C_t/C_o) \text{ W/sq.m.} \dots(4)$$

⁹ A similar equation is for combustion of petroleum octane: $2C_8H_{18} (l) + 25O_2 (g) \rightarrow 16CO_2 (g) + 18H_2O (l)$, where *l* and *g* are litres and grams respectively, and the value for the energy produced, -11020 kJ / mol, tells us that for every one of these reactions (with inputs of 2 gram mol weights(GMW) of octane, 25 GMW of oxygen, and outputs of 16 GMW of carbon dioxide, and 18 GMW of water), 11,020 kilojoules of heat is released, rather more than is attributable to the radiative forcing by 16 GMW of CO₂. The words hydrocarbon and combustion do not appear in the Index in the IPCC's Solomon *et al.* 2007. For global estimates of net heat emission from energy production, see Nordell and Gervet (2009).

¹⁰ For a full account of the photosynthesis process, see Farquhar *et al.* 1980.

¹¹ This account is based on the article *Radiative Forcing* in Wikipedia, accessed 14th March 2011. See also Ramaswamy *et al.* (TAR, 2001:354).

where C_t is the current or future level of atmospheric carbon dioxide and C_o is the pre-industrial level (280 ppm), and F is measured in Watts per square metre. A typical value of λ is said to be 0.8 K/W/m², which gives a warming of 3 °C for a doubling of CO₂ (from 280 ppm to 560 ppm). Equations (3) and (4) disclose no physical relationships, and equation (4) (derived by Myhre *et al.* 1998, see also Ramaswamy *et al.* 2001:361) also describes no *process* whereby changes in the ratio of the present or future levels of [CO₂] to the base level of 280 ppm generate radiative forcing. Moreover the parameters λ and 5.35 are not data-based, e.g. by use of linear least squares regression, being set in models only *ex post* at whatever level delivers some sort of match to the observations (see Myhre *et al.* 1998). Clearly the only *variables* in (3) and (4) are the base and projected levels of [CO₂]. But Myhre's equation does have the property that because it takes the *ratio* of current [CO₂] to the base year (1750) level, the radiative forcing F inexorably *increases* along with the rising level of [CO₂]. This is contrary to the Arrhenius formulation using only increases in absolute levels of [CO₂]. In his words "if the quantity of carbonic acid [i.e. CO₂] increases in geometric progression, the augmentation of the temperature will increase nearly in arithmetic progression" (1896:267 and Table VII), i.e. less than in proportion to the increase in [CO₂].

As a result, this model consistently overstates its forecast of global temperature from 1959 to 2010 by 0.0112oC p.a. (0.112°C per decade), and higher by 0.59°C for 2010 compared with the Gistemp global temperature of 14.83°C in that year. That is not a trivial difference, and it gets worse over time, especially if the claimed forcings of all the other greenhouse gases are added to that from [CO₂].

Thus the equations (3) and (4) may look impressive but they lack the precision of chemical formulae like (1). Climate scientists should at least attempt to evaluate an equation like:

$$\Delta T_t = a + b(\text{CO}_2)_t + c(\text{H}_2\text{O})_t + u_t \dots(5)$$

Here H₂O includes *both* the water vapour generated in the combustion process *and* the steam produced by the generation of electricity. There is no consideration of such a formula in the science academy texts relied on by Garnaut's SCC (e.g. Allison 2010, Royal Society 2010). Climate scientists have never attempted to provide statistically significant values for the coefficients b and c . Instead, the IPCC's Solomon *et al.* (2007) specifically excludes consideration of any of the (H₂O) outputs of hydrocarbon combustion and power generation, which means its implied derivation of ΔT_t bears no relation to the multi-variable combustion output and power formula (5).

However, it is to be noted that equation (5) does not lend itself to econometric analysis, because the CO₂ and H₂O variables are not independent of each other, as they are co-determined during the combustion process and power generation processes. This means that the implied correlations of temperature anomalies and increasing [CO₂] all *unavoidably* include the water vapour component as an ingredient of the radiative forcing in the IPCC's Solomon *et al.* (2007) and the rest. But it is perfectly possible to generate statistics of not just emissions of CO₂ but also of emissions of water vapour and steam in the processes of hydrocarbon combustion and power generation.

Kelly (2010:630) notes that in the various applicable chapters in Solomon *et al.* (2007), there is no mention that most (but not all) anthropogenic water vapour is in the form of steam. That steam is over and above the flue exhaust H₂O in the combustion process. For example, the condensate flow rate of the water used to produce steam by the heating of the combustion process (see equation 1 above) at full load in a 500 MW plant is about 6,000 US gallons per minute (400 L/s) (“Thermal Power Station”, *Wikipedia*, accessed 14 April 2011), or c.12.6 million tonnes p.a.¹² Most of the electricity in the United States is produced using steam turbines. Coal is the most common fuel for generating electricity in the United States. In 2009, 45% of the USA’s nearly 4 trillion kWh of electricity used coal as its source of energy, but gas, oil, and nuclear power stations also mainly produce steam which is of course used to drive the generators producing their electricity.

Global energy use was 98,022 TWh in 2009 (*Wikipedia*, accessed 14 April 2011). That would have had an associated generation of steam of 303.75 GtH₂O (author’s calculations), very much larger (ten times) than the gross global output of CO₂ from hydrocarbon combustion of 30 GtCO₂ in 2009 (le Quéré *et al* 2009) – and of course a very much larger radiative forcing. Thus in the language of the Garnaut *Reviews*, it is water, not carbon dioxide, that is the “pollutant”. As I show below, this is confirmed by multivariate econometric analysis which is fully capable of generating those coefficients from data on levels of [CO₂] and [H₂O] at specific locations where these variables are spatially independent of each other, and reveals that *b* is usually close to zero, even negative, while *c* is large and highly statistically significant.

Climate scientists are aware of the Clausius-Clapeyron relation that defines the maximum partial pressure of water vapour that can be present in a volume of atmosphere in thermodynamic equilibrium as a strongly increasing function of temperature, but have failed to quantify it accurately in the context of their predicted climate change. This maximum is known as the saturation vapour pressure, *e_s*:

$$e_s(T) = e_s(T_0) \exp^{-L/Rv(1/T - 1/T_0)} \dots (6)$$

“where *L* is the latent heat of the appropriate phase transition (vapour to liquid at warm temperatures, vapour to solid at sufficiently cold temperatures), *Rv* is the gas constant for water vapour, and *T₀* is a reference temperature. At the freezing point, *e_s* is 614 Pa or 6.14 mb; *L/Rv* = 5419 K for condensation into liquid and 6148 K for condensation into ice” (Pierrehumbert *et al.* 2007:145). While this expression shows that the ability of the atmosphere to hold water vapour rises with temperature, it does not imply it is changes in *T* that increase both the *amount* of that water vapour and the saturation level *e_s*.

¹² “Typical water consumption for a 1000 MWe plant - providing electricity to perhaps 1 million people in an industrial country - might be 75 megalitres per day, or the equivalent of 25 Olympic-sized swimming pools. This equates to about 0.05% of the average flow rate of the Rhine, Rhone and Danube Rivers”, World Nuclear Association, nd. This implies that the USA and EU have about 500 1000MWe-equivalent power stations, requiring 25% of the flow of the Rhine, Rhone, and Danube rivers.

Thus while satellite observations indicate that the response of global evaporation to rising temperature is close to that implied by (6), with the cooling effect entailed by evaporation and associated precipitation, the GCM models deployed by climate scientists apparently *reduce* the evaporation implied by Clausius-Clapeyron, presumably because that increases potential global warming (Kininmonth 2010:61). Hence Pierrehumbert *et al.* caution against what they term “wholly indefensible statements which simply invoke the Clausius-Clapeyron relation...[as being] at the root of the behaviour of water vapour, *but the proper use of the relation hinges on identifying the temperature to which the relation should be applied; it's not the surface temperature, and the effect of the relation on evaporation is of little relevance to water vapour feedback*” (2007:180, my emphasis).

As a result of not acting on the advice of Pierrehumbert *et al.* (2007), Solomon *et al.* (2007) wrongly attribute increases in water vapour only to rising surface temperatures, although using those increases to raise the radiative forcing they ascribe to a doubling of [CO₂] is enough for them to triple the temperature change due to that by 2100 from 1°C to over 3°C. This artefact is termed the “positive feedback” effect whereby rising temperature raises evaporation whose own radiative forcing would be much stronger than that of atmospheric carbon dioxide alone, were it not for precipitation rising *pari passu* with evaporation. But Randall and Wood *et al.* (2007, in Solomon *et al.* 2007:633) and Garnaut (2008:81 and 2011:26-27) downplay the role of precipitation in cancelling out rising evaporation, instead they often predict increasing drought in various regions despite the rising water vapour they rely on for the infinitely spiralling further temperature increases that deliver their “runaway” greenhouse effect.

In addition, Solomon *et al.* (2007) appear to believe that the positive feedback effect of rising temperature on the level of atmospheric water vapour outweighs that of direct solar radiation. However, as Kiehl and Trenberth show (e.g. 1997:206), incoming solar radiation of 342 W/sq.m. generates “evapotranspiration” of 78 W/sq.m., which is 0.228 W/sq.m. of incoming solar radiation, whereas the evapotranspiration of total radiative forcing from long lived greenhouse gases, estimated at 2.63 W/sq.m. in 2005 by Forster and Ramaswamy *et al.* (2007, in Solomon *et al.*, 2007:141), is on that basis only 0.599 W/sq.m., less than 1% of that from solar. Those authors do not mention evapotranspiration arising from solar radiation, but assert that the derisory increase from the GHG forcing of only 2.63 W/sq.m. is enough to raise the increase in global temperature from rising concentrations of greenhouse gases by a factor of at least two.

However, Pierrehumbert *et al.* also assume that increases in atmospheric water vapour [H₂O] stem only from rises in temperature, and are independent of processes like combustion of hydrocarbons that directly release water vapour into the atmosphere. Such releases *must* be just as potent *pro rata* – in terms of generating radiative forcing by reducing outgoing long wave radiation (OLR) – as increases in [H₂O] resulting from rising temperature. Moreover, like those increases in [H₂O], the increases produced by combustion must also be “approximately logarithmic in specific humidity once the [total] concentration is sufficiently large to saturate the principal absorption bands” (*ibid.*). They also find that each doubling of water vapour reduces OLR by about 6W/m², and add:

This is about 50% greater than the sensitivity of OLR to CO₂. The idea that small quantities of water vapour can have a lot of leverage in climate change has a fairly long history, and is now widely recognized. Water vapour feedback was included in the very first quantitative calculations of CO₂-induced warming by Arrhenius [1896], and the importance of water vapour aloft was implicit in such calculations.... In fact, at each latitude the greenhouse effect of water vapour is approximately twice that of CO₂ (Pierrehumbert *et al.* 2007:146, 148).

For example, we know that in 2008-2009 combustion of hydrocarbon fuels generated emissions of about 8.6 billion tonnes of carbon, which equates to 31.4 billion tonnes of CO₂, (i.e. 31.4 GtCO₂) (le Quéré *et al.*, 2010) – and using data in Gaffen and Ross (1999) this implies direct water vapour emissions (just from the combustion process of burning a hydrocarbon fuel using oxygen) of 17.5 GtH₂O.¹³ But using the first Pierrehumbert *et al.* figure above, the radiative forcing (RF) from this addition to [H₂O] is 50 per cent higher than that of increased atmospheric CO₂.¹⁴ According to the IPCC (Forster and Ramaswamy 2007: 141), the radiative forcing per GtCO₂ is 0.0019 Watts/sq.metre, so that from changes in [H₂O] is 0.0028 W/sq.metre. That means hydrocarbon combustion alone (excluding steam) generated total RF of 0.0047 W/sq.m. *per tonne* of H₂O and CO₂ in 2008-09. Thus total RF from just hydrocarbon combustion across the whole surface of the globe was 5.09E+14 Watts, of which 45 per cent was the contribution of the water vapour from fossil fuel combustion. Yet Garnaut's SCC never notices this effect, and it is dismissed as of no consequence by the IPCC's Forster and Ramaswamy *et al.* (2007:185).¹⁵

¹³ "Based on carbon emissions data (Marland *et al* 1994) we estimate global water vapor emission from fossil fuel consumption to be of order 10¹² (in 1960) to 10¹³ (in 1990) kg yr⁻¹", Gaffen and Ross, 1999. Carbon emissions were 5.57 GtC in 1990, yielding a factor of 5.57/10 for the ratio between emissions of water vapour and carbon, in Giga tonnes, from hydrocarbon combustion, namely 1.795, say 1.8 (or 0.49 Gt H₂O to 1 Gt of CO₂).

¹⁴ "Specifically, the team found that if Earth warms by 1.8 degrees Fahrenheit, the associated increase in water vapor will trap an extra 2 Watts of energy per square meter (about 11 square feet). That number may not sound like much, but add up all of that energy over the entire Earth surface and you find that water vapor is trapping a lot of energy... we now think the water vapor feedback is extraordinarily strong, capable of doubling the warming due to carbon dioxide alone." Water Vapor Confirmed As Major Player In Climate Change. *ScienceDaily*. NASA/Goddard Space Flight Center (2008, November 18). This comment refers only to the claimed water vapour "feedback" and ignores all anthropogenic additions to [H₂O] from the steam and direct emissions of water vapour during the combustion and power generation processes.

¹⁵ Forster and Ramaswamy *et al* state "The emission from water vapour from fossil fuel combustion is significantly lower than the emission from changes in land use (Boucher *et al.*

The implications are profound, because clearly the RF and resulting global warming from hydrocarbon fuel combustion must be seriously underestimated by disregarding the water vapour component of the total output. However Lim and Roderick (2009) have shown that in reality, while there is some evidence for increasing evaporation between 1970 and 1999, it has been *exactly* matched by increased rainfall. Thus just as the IPCC and Garnaut's *Reviews* generally assume reducing uptakes of CO₂ emissions by the biospheres, they also ignore that rising [H₂O] is matched by rainfall, preferring instead to stress the claimed but as yet undemonstrated positive feedback effect of rising temperature through evaporation to yet higher temperatures. This explains why they always ignore the positive social benefits of increases in both evaporation and rainfall, and implicitly treat the resulting increases in agricultural productivity as negative social "externalities".

5. Previous Econometric Analysis of Climate Change

There have been surprisingly few attempts to use econometric analysis to separate out the proximate causes of climate change.

Few researchers have used time domain econometrics methods to analyze climate change. Apart from Kaufmann and Stern (1997) and Stern and Kaufmann (1999), only Tol and de Vos (1993, 1998), [and] Tol (1994), explicitly use econometric time series methods to investigate the causes of climate change... (Stern and Kaufmann 2000:412).¹⁶

Kaufmann *et al.* (2006b) made further use of econometric methods, and comment:

...statistical models of the relationship between surface temperature and radiative forcing that are estimated from the observational temperature record often are viewed skeptically by climate modelers. One reason is uncertainty about what statistical models measure. Because statistical models do not represent physical linkages directly, it is difficult to assess the time scale associated with statistical estimates for the effect of a doubling in CO₂ on surface temperature...The results of this analysis indicate that the statistical methodology used by Kaufmann *et al.* (2006a) to analyze the instrumental temperature record is able to provide an accurate and unbiased estimate for the true (model specific) estimate for the [transient climate response] TCR. This accuracy gives us added confidence in the statistical estimates for the TCR that are estimated from the instrumental temperature record. Although the transient climate response does not include the full effects of deep ocean mixing and feedbacks, it does represent the challenge for the twenty-first century and therefore is important for attributing climate

2004)" (2007:185). In reality, Boucher *et al.* "estimate a global mean radiative forcing in the range of 0.03 to +0.1 Wm⁻² due to the increase in water vapour from Irrigation" (2004:597), and their lower figure is effectively the same as mine (0.028 W/sq.m.) for that from combustion alone (ignoring the much larger contribution from steam emitted during power generation).

¹⁶ Tol and Vellinga (1998) used econometric analysis to separate the enhanced greenhouse effect from the influence of the sun at the top of the atmosphere (TOA), while Tol and de Vos (1998) used Bayesian analysis. Neither paper considers the role of atmospheric water vapour.

change to human activity and efforts to formulate economically efficient policy.

Hegerl and Zwiers *et al.* (2007, in AR4, Solomon *et al.* 2007) failed to respond to that challenge. Although claiming they would attempt to “understand climate changes that result from anthropogenic and natural external forcings” (p.667), they never report any regression results estimating the relative values of those forcings. They concede (p.668) that “attribution studies ... assess whether the response to a key forcing, such as greenhouse gas increases is distinguishable from that due to other forcings (Appendix 9A). These questions are typically investigated using a multiple regression of observations onto several fingerprints representing climate responses to different forcings... see Section 9.2.2”.

There is no trace of the results of any such analysis anywhere in Hegerl and Zwiers *et al.* 2007, least of all in either of their referenced Section 9.2.2 or their Appendix 9A. The latter (pp.744-745) does have a textbook account of multivariate regression, but reports no results at all. Yet it is Hegerl and Zwiers *et al.* (p.666) who so confidently assert “greenhouse gas forcing has *very likely* caused *most* of the global warming over the last 50 years” (our emphasis), where “very likely” means “more than 90 percent probability” (Solomon *et al.* 2007:121) and “most” must mean more than 50 percent where only two independent variables are considered. Had these authors done some regression analysis they could have been more precise, but they never did, nor do they report any by others.

6. Regression analysis of climate science counterfactuals

Using the Mauna Loa data set on additions to atmospheric CO₂ from 1958 to 2009, and regressing Gistemp temperature anomalies against it, we have these results:

Table 1 Regression of Gistemp GMT anomalies (oC) against Radiative Forcing of [CO₂] (W/sq.m) 1959-2010.

| Predictors of Anomalies | t-statistics | p-values |
|--|--------------|----------|
| Constant | -8.260 | 0.000 |
| Radiative Forcing (GtCO ₂) | 12.749 | 0.000 |
| Adjusted R ² | 0.76 | |
| Standard Error | 13.97 | |
| Durbin-Watson | 1.31 | |

Sources: NASA-GISS; NOAA-CDIAC; Solomon *et al.* 2007:141.

These results look impressive, with a high R² of 0.76 and a statistically significant coefficient on the radiative forcing of the [CO₂] variable (i.e. atmospheric CO₂ as measured at the Mauna Loa Observatory). But as von Storch and Zwiers note in their textbook (1999), comparison of time series using least squares regression has a high risk of serial or spurious correlation, and the Durbin-Watson statistic at 1.313, which is well below the benchmark 2.0, shows that to be the case in this instance.

This problem is usually addressed by taking year-on-year differences in the data (e.g. Breusch and Vahid, 2011), with the results shown in Table 2, which may explain why the IPCC (Solomon *et al.* 2007) and Garnaut (2008 and 2011) do not report regression results, for while the Durbin-Watson statistic at 2.85 rules out serial correlation, the R² falls to .04, while the CO₂ variable is totally statistically insignificant. This means there is no basis for the claim by Solomon *et al.* (2007:10) that changes in anthropogenic atmospheric greenhouse gases account for “most” (i.e. more than 50 per cent) of observed temperature change since 1958, as claimed by Garnaut (2008,2011), Stern (2006), and Hegerl and Zwiers *et al.* (2007:670).

Table 2 Regression of changes in surface GMT on changes in [CO₂] Radiative Forcing

| Predictors of Changes in Anomalies | t-statistics | p-values |
|------------------------------------|--------------|----------|
| Constant | .095 | 0.925 |
| Changes in RF (GtCO ₂) | 0.455 | 0.651 |
| Adjusted R ² | -0.016 | |
| Standard Error | 16.156 | |
| Durbin-Watson | 2.846 | |

Sources: NASA-GISS; NOAA-CDIAC; Solomon *et al.* 2007:141.

Garnaut’s SCC overlooks that Hegerl and Zwiers as lead authors of the crucial Chapter 9 (“Understanding and Attributing Climate Change”) in IPCC’s AR4 (Solomon *et al.* 2007) must have been well aware that standard multivariate regression analysis of changes in temperature as a function of changes in atmospheric CO₂ and natural influences (e.g. water vapour) does not yield their desired outcomes, otherwise how can we explain the absence of any reporting of such analysis by Hegerl and Zwiers *et al.* 2007 when Zwiers is also co-author of the standard text (1999) on statistical analysis of climate change? Instead Hegerl and Zwiers *et al.* (2007) resort to developing a new method of analysis they have named “detection and attribution of fingerprints”. But this system does not appear in standard textbooks, as it twists the raw data into their desired shapes by use of modelled parameters rather than using only raw data. Thus although their model has the form $y = Xa + u$ where y is the “filtered” climate variable and X is a matrix of independent variables that could determine y , while a is a vector of scaling factors, in their hands X , the matrix for the independent variables, contains only “signals that are estimated” by climate models, rather than real world data like that I have presented above (Hegerl and Zwiers *et al.* 2007: 744).

7. Counterfactuals, Carbon Dioxide, and Climate Change

I gave above a brief account of John Snow’s counterfactual analysis of cholera by collecting statistics on the distribution of sources of water supply by households and of deaths from cholera. The problem with the bivariate regressions of global temperature change and atmospheric carbon dioxide reported in the previous section is that the only independent variable is changes in atmospheric CO₂. In that regard such regressions are like those implicitly relied on by the medical establishment in London in 1854, which could as easily have proved that the increasing incidence of cholera outbreaks was due to the equally well

attested increase in use of horse-drawn vehicles, omnibuses, cabs, and the carriages of the rich, with horses' exhalation a plausible source of the miasma passionately believed by the Garnauts of the day to be the cause of cholera.

The modern equivalent of John Snow's careful mapping of cholera deaths and water supply is climate data sets from over 1200 locations produced by the US Government's NOAA that are, if not deliberately ignored, certainly overlooked by the climate change establishment, perhaps because they include time series of a wide range of climatic variables, not just temperature.

These location-specific and multivariate data sets provide counterfactuals simply because although atmospheric carbon dioxide is known to be a "well-mixed" greenhouse gas (WMGHG), like London's miasma in 1854, which means it is at much the same level everywhere, from Point Barrow in Alaska to Cape Grim in Tasmania to Mauna Loa in Hawaii, temperature *trends* are NOT the same everywhere, and are in general more closely associated with other climatic variables, such as the amount of sunlight reaching the surface, and, above all, the level of atmospheric water vapour, i.e. [H₂O]. My regressions reported in Tables 3, 4, and 5 analyse the trends in annual mean maximum or minimum temperatures at Point Barrow in Alaska, Hilo (near Mauna Loa) in Hawaii, and New York for the period 1960-2006. The respective trends differ considerably, although the R² correlation coefficients do not suggest good fits for linear rather than polynomial trends. The *nil* linear trend in the annual mean temperature at Mauna Loa itself from 1978-2006 with R² of 0.1 contrasts with that for atmospheric carbon dioxide (R² 0.99), and in part explains the poor correlation in Table 4.

The regression analysis (Table 3) of the NOAA climate data for Pt Barrow for 1960 to 2006 (when this data set terminates) provides a striking example of John Snow's counterfactual analysis, as it shows that changes in the level of atmospheric carbon dioxide play no role at all in explanation of the significant warming trend in minimum temperatures, while the role of atmospheric water vapour [H₂O] is strong and very highly significant (99%), along with opacity of the sky at night (OPQ). Similar results are found for various locations in Hawaii, such as Hilo (Table 4), and New York City (Table 5, and Fig.3), and many others. The analysis for Hilo reported in Table 4 has changes in average annual maximum temperatures as the independent variable, with changes in solar radiation at ground level ("Avglo") replacing the sky opacity variable, as clearly solar radiation is more relevant to maximum than minimum temperatures.

Table 3 Regression of climate data, Point Barrow 1960-2006

| Change in Mean Minimum Temperatures | t-statistics | p-values |
|-------------------------------------|--------------|----------|
| Constant | -1.12 | 0.91 |
| Change in [CO ₂] | 0.06 | 0.95 |
| Change in [H ₂ O] | 4.96 | 0.000 |
| Change in OPQ | 4.19 | 0.000 |
| Standard Error | 1.04 | |

| | |
|-------------------------|--------------|
| Adjusted R ² | 0.59 |
| Durbin-Watson | 2.846 |

Sources: <http://rredc.nrel.gov/solar/old-data/nsrdb/1961-90/dsf/data> and <http://rredc.nrel.gov/solar/old-data/nsrdb/1991-2005/statistics/data> .

Note: High *p*-values with low *t*-values indicate very low statistical significance of [CO₂], and conversely for high *t* and low *p* values for [H₂O] and sky opacity (OPQ) coefficients.

Table 4 Multivariate regression of climatic variables, Hilo (Hawaii) 1960-2006

| Change in Annual Mean Maximum Temperatures | t-statistics | p-values |
|--|--------------|----------|
| Constant | -1.13 | 0.264 |
| Change in [CO ₂] | 1.35 | 0.186 |
| Change in [H ₂ O] | 4.67 | 0.000 |
| Change in Avglo | 3.72 | 0.000 |
| Standard Error | 0.565 | |
| Adjusted R ² | 0.356 | |
| Durbin-Watson | 2.834 | |

Sources: see Table 3.

Note: Independent variables in Table 4: ΔCO_2 = annual changes in atmospheric carbon dioxide at Mauna Loa Slope Observatory; ΔAvglo = change in annual average daily *in situ* total solar radiation, i.e. sum of direct and diffuse radiation less albedo (Wh/m²); $\Delta\text{H}_2\text{O}$ = annual changes in precipitable water vapour (cm).

Table 5 Regression of climate data, New York City (JFK) 1960-2006


| Change in Annual Mean Maximum Temperatures | t-statistics | p-values |
|--|--------------|----------|
| Constant | -0.39 | 0.70 |
| Change in [CO ₂] | 0.14 | 0.89 |
| Change in [H ₂ O] | 9.64 | 0.000 |
| Change in OPQ | -1.49 | 0.14 |
| Standard Error | 1.86 | |
| Adjusted R ² | 0.69 | |
| Durbin-Watson | 2.692 | |

Sources: see Table 3.

Note: The independent variables in Table 4 are: ΔCO_2 = annual changes in atmospheric carbon dioxide at Mauna Loa Slope Observatory; ΔOPQ = changes in annual average opacity of the sky; $\Delta\text{H}_2\text{O}$ = annual changes in precipitable water vapour (cm).

These counterfactuals all show no statistically significant correlations between atmospheric carbon dioxide levels (i.e. $[CO_2]$) and changes in temperature, and thereby fully support the conclusions of Paulo Cesar Soares (2010:111), that there is no causal relation between $[CO_2]$ with global warming, given the absence of evidence for changes in $[CO_2]$ preceding temperature either for global or local changes, and that the greenhouse effect of $[CO_2]$ is very small compared with that of water vapour.

Fig.3 Annual changes in mean temperature, $[CO_2]$ and $[H_2O]$, New York 1960-2006



Sources: see Table 3.

Note: This graph explains the regressions reported in Table 5, showing why the annual changes in atmospheric carbon dioxide concentrations do not explain temperature variability in New York, unlike changes in atmospheric water vapour.

8. Water Vapour and Climate Change

Projections of global surface warming as large as $5.8^{\circ}C$ over the next 100 years are based not on the direct contribution to atmospheric water vapour $[H_2O]$ from combustion of hydrocarbons but on increases in $[H_2O]$ arising from rising temperatures attributed to increasing $[CO_2]$, hence the term “feedback” to describe a second round impact of increasing $[H_2O]$ on temperature (Minschwaner and Dessler 2004, Dessler *et al.* 2008). In most global climate models, an initial warming caused by additional CO_2 and other greenhouse gases leads to enhanced evaporation at the surface and higher specific humidity of the

atmosphere. Since water vapour is a strong infrared absorber (Arrhenius 1896), the added moisture causes further warming. But how much? Minschwaner and Dessler (2004:1281) estimate the change in water vapour in the Upper Troposphere (near 215 mb pressure or altitude of about 12 km.) is between 8.5 and 9.5 ppmv per 1°C increase in surface temperature. That implies a total increase in [H₂O] from the 0.7°C rise in global surface temperature between 1900 and 2000 of just 5.95-6.65 ppmv., to be compared with the apparent increase in [CO₂] from 1900 to 2000 of 91 ppmv. Using that calculation, it is difficult to see how Minschwaner and Dessler (2004:1272) arrived at “projections of surface warmings as large as 5.8°C over the next 100 years” because of the “positive water vapour feedback in the atmosphere”.

However, Dessler *et al.* (2008) claim to have demonstrated a substantial water-vapour climate feedback by comparing satellite data on tropospheric specific humidity q in the El Niño of December 2006-February 2007 with that in the cooler La Niña of December 2007-February 2008, with q higher in the former. Their paper does not explain why it was cooler in the La Niña despite higher [CO₂] than a year before, and shows no feedback impact on temperature in La Niña from the higher q during El Niño.

Paltridge *et al.* (2009) found that specific humidity in the National Centres for Environmental Prediction/National Centre for Atmospheric Research (NCEP/NCAR) reanalysis declined between 1973 and 2007, particularly in the tropical mid and upper troposphere, the region that plays the key role in the water vapour feedback. They pointed out that climate models tend to maintain constant relative humidity at each atmospheric level, and therefore have an increasing absolute humidity at each level as the surface and atmospheric temperatures increase. This behaviour in the upper levels of the models produces a positive feedback which more than doubles the temperature rise calculated to be the consequence of increasing atmospheric CO₂. Total water vapour in the atmosphere may increase as the temperature of the surface rises, but if at the same time the mid- to upper-level concentration decreases then water vapour feedback will be negative. Minschwaner and Dessler show exactly that effect (2004: Figs. 1 and 2).

These suggestions by Paltridge *et al.* (2009) have been contested by Dessler and Davis (2010) who conceded that “if borne out, this result suggests potential problems in the consensus view of a positive water vapour feedback”, but produce further reanalysis rejecting the findings of Paltridge *et al.* However, Hansen *et al.* (2011) note that “the latent energy associated with increasing atmospheric water vapour in a warmer atmosphere is an order of magnitude too small to provide an explanation for the high estimates of atmospheric heat gain” (like those in Dessler *et al.* 2008), and thereby confirms the findings of this chapter, that the direct anthropogenic contributions to [H₂O] of currently over 300 GtCO₂ p.a. are much more significant than the claimed positive [H₂O] feedback effects from the essentially trivial increase in global temperature since 1900 of about 0.007°C a year.

9. The Benefits of Hydrocarbon Fuel Combustion Emissions

Garnaut's first Update paper (2011) has the title "Weighing the costs and benefits of climate change action", and like the fifth is in breach of ASX disclosure rules, because at no point does it mention the undoubted benefits of hydrocarbon combustion in terms of (1) the fertilization effect of elevated atmospheric carbon dioxide on primary production, and (2) the benefits in terms of enhanced crop yields of higher global rainfall. For Garnaut (2008 and 2011), hydrocarbon combustion produces only costs, in the form of as yet unproven large rises in global mean temperatures (even though they would also have benefits in the form of higher crop yields and longer growing seasons, especially in all higher latitude zones), and the benefits of restricting use of hydrocarbons consist *only* of avoidance of the supposed costs of rising temperatures.

Ironically, there is a contradiction between the SCC's basic position that hydrocarbon emissions produce only costs, and that the biospheric sinks are close to saturation, as claimed by Raupach *et al.* (2008, etc.) while in both his 2008 and 2011 *Reviews*, Garnaut devotes space to the role of expanding agro-forestry in order to reduce atmospheric carbon dioxide concentrations through their absorption of substantial volumes of those emissions, which would not be possible if Raupach *et al.* were correct. Yet the 2008 *Review* raises the historic growth rate of the concentration of CO_{2e} at 0.29 per cent p.a. (1958-2010) to his projection of 1.0137 per cent p.a. from 2010 to 2100 by suppressing the known take-up of more than half of carbon dioxide emissions by the globe's biospheres (Knorr 2009). This is justified in section 2.3.2 (in SCC) on the carbon cycle which at face value seems unexceptionable but nevertheless displays palpable bias. The first is the stress on the apparently fast growth of fossil fuel emissions between 2000 and 2008 of 3.4 per cent p.a., without mentioning that in that period the atmospheric concentration of CO₂ increased by only 0.296 per cent p.a.

This distinction between the respective growth rates of emissions and of the atmospheric concentration is very important but never mentioned in either the 2008 or 2011 *Reviews*, and may betray some ignorance of dependence of percentages on respective base levels. Total annual emissions of human-generated carbon dioxide are approaching 10 billion tonnes of carbon-equivalent CO₂ (i.e. GtC), but the absolute atmospheric concentration is nearly 830 GtC. An increase of three percent in the first figure, of which 55 percent is absorbed by the biospheres, becomes just 0.016 per cent of the second figure.

At the observed growth rate of [CO₂] of 0.295 per cent p.a. over the longer period from 1959 to 2009, it will take until **2134** for the so-called pre-industrial level of the atmospheric concentration of 280 ppm to double to 560 ppm; and the 2009 level of 387 ppm will not double until **2244**. The *Reviews'* portrayal of the extreme urgency of preventing these gentle increases is not compelling, unless one accepts their projection of 1565 ppm CO_{2e} by 2100.

The SCC justifies its alarmism by claiming that the strength of the carbon sinks (which have on average absorbed 56 per cent of carbon dioxide emissions since 1958) has been "declining" over the last five decades, which if true would result in an increasing airborne fraction of emissions from the present average level of 44 per cent p.a. It offers no statistical evidence beyond its repeated citations of papers by J.P. Canadell, C. Le Quéré, P. Fraser, and M. Raupach, of which none of those cited in SCC section 2.3.2 has been peer-reviewed. Instead, as already noted above, with another omission of material facts, the 2011 *Review*

ignores the peer-reviewed paper by W. Knorr (2009) in a mainstream journal, which concludes “despite the predictions of coupled climate-carbon cycle models, no trend in the airborne fraction can be found”.

Another peer-reviewed paper (Curtin 2009) that appeared in the same month as Knorr’s provided substantive statistical support for the benefits conferred on mankind by the rising level of atmospheric carbon dioxide through its enhancement of global cereal production based on the increasing photosynthesis - on which all living matter is wholly dependent - facilitated by increases in [CO₂]. That “externality” is never mentioned in Garnaut 2008 and 2011 even when admitting that expanding agro-forestry can increase the uptakes of carbon dioxide emissions that elsewhere he projects as falling to nil (2008: Fig.2.7). The role of rising temperature and CO₂ in raising photosynthesis - and thereby yields - is shown in Fig.4. Evidently the more CO₂ is available at any given temperature the higher is the net primary productivity (NPP). Similarly, the higher the temperature at any level of CO₂, the higher is the NPP (left axis) up to the limits specified. Combining higher temperature with higher CO₂ again leads to higher NPP. We are now at 390 ppm and <15°C, so have some way to reach the optimum implied by the feasible levels of 650 ppm and leaf temperature of 30°C.

Similarly, here Table 6 confirms the concurrence in Fig.4 of rising atmospheric carbon dioxide with rising terrestrial uptakes of carbon dioxide and rising world cereals production both in total and per capita. The *Reviews* will incur a heavy burden of responsibility, if their proposals to secure a reduction of carbon dioxide emissions succeed in reducing the atmospheric concentration to levels like that shown for 350 ppm in Fig.4 that are unlikely to support the food requirements of a world population of nearly 7 billion people now, and possibly 9 billion by 2050.²²

Table 6 Carbon Dioxide and World Cereals Production 1961-2007

| | 1961 | 2007 | % Change |
|--|------|------|-------------|
| Atmospheric concentration of CO ₂ (ppm) | 673 | 815 | 121 |
| Anthropogenic emissions of CO ₂ (GtC) | 4.06 | 9.94 | 245 |
| Total terrestrial uptakes (GtC) | 0.9 | 2.98 | 331 |
| Total cereals production, Gt | 0.88 | 2.34 | 265 |
| Carbon content (average 40%) GtC | 0.35 | 0.94 | 268 |
| World population (billions) | 3.08 | 6.6 | 214 |


²² “The fundamental reason why carbon dioxide in the atmosphere is critically important to biology is that there is so little of it. A field of corn growing in full sunlight in the middle of the day uses up all the carbon dioxide within a meter of the ground in about five minutes. If the air were not constantly stirred by convection currents and winds, the corn would stop growing” (Dyson 2007).

| | | | |
|---|-------------|-------------|------------|
| Cereal production per capita, tonnes | 0.28 | 0.35 | 125 |
| Carbon content per capita, 40% average, tonnes | 0.11 | 0.14 | 127 |

Source: Le Quéré, Canadell *et al.* 2009, www.globalcarbonproject.org, FAO 2009.

Note: The world's uptakes of atmospheric carbon dioxide are not limited to cereals - other food and fruit crops, livestock, forestry, and fisheries are all absorbers through photosynthesis.

Fig. 4 The response of light-saturated net photosynthesis (Pn) to changes in leaf temperature at different atmospheric CO₂ concentrations.



Sources: Norby and Luo (2004) and Long (1991), with permission of Blackwell Science.

Note: Arrows indicate the shift in optimum temperature with increasing CO₂ concentration.

10. Conclusion

This chapter has produced statistical and econometric analysis showing that the mainstream science of climate change has been too focussed on a single supposed causative agent, carbon dioxide emissions (about 30 GtCO₂ p.a.), to the serious neglect of the much larger volumes of anthropogenic water vapour produced by the combustion of hydrocarbon fuels, both by direct creation of water vapour in the combustion process (18 GtH₂O p.a.), and by the much larger volume of steam created by the power generation process (300 GtH₂O). It is true that individual injections of water vapour to the atmosphere have a short residence time there before descending as precipitation (Kelly, 2010), but increases in the average level of [H₂O] over a year remain significant.

The implications of full accounting for the radiative forcing attributable to anthropogenic water vapour are confounding. On the one hand, it is conceivable that it has been unwittingly included in the radiative forcing ascribed to rising [CO_{2e}], while on the other it may be an addition to that forcing, which would mean that temperature projections are being understated. Alternatively, it may be that the forcing ascribed to positive feedbacks from rising evaporation due to the observed annual global temperature increase of 0.007°C p.a., which raises the IPCC's projected temperature increase from a doubling of [CO_{2e}] to 3°C from the 1°C due to [CO_{2e}] alone, despite lack of evidence for this effect, should be reassigned to anthropogenic water vapour.

Ross Garnaut ends his *The Science of Climate Change* with this comment (p.44):

Scepticism is an essential part of the scientific process and serves to move the science towards greater understanding and agreement. Once a theory is put forward by a researcher, it is discussed, analysed and criticised by the wider scientific community. Further tests, modelling and research then respond to that questioning. The consequent exchanges determine whether the initial conclusions hold, need refining or are rejected.

I offer this critique in that same spirit (as in Curtin 2009a). In particular, this chapter's econometric regressions of temperature changes at key locations on a range of climatic variables, including especially atmospheric water vapour, show that atmospheric carbon dioxide plays a negligible role. Exceptions do not prove a general rule, especially when the exceptions are many hundreds of locations across the length and breadth of the USA, still either the worlds' largest or second largest emitter of carbon dioxide.

It would be a good start if the Australian government (and all governments sponsoring the IPCC's Fifth Assessment Report (due in 2013/14) referred in future more accurately to its proposed new impost as a "Tax on Plant Growth and Rainfall", given the data here showing (1) that the respective radiative forcings from emissions of water vapour and carbon dioxide during hydrocarbon combustion and power generation are much more potent from the former than the latter, and (2) that rises in temperatures, levels of atmospheric carbon dioxide, and atmospheric water vapour have all been consistent with the increases in per capita cereals production since 1960.

Acknowledgments

I am particularly indebted to Anthony Kelly, and also to Anthony Cox, Alan Emmerson, Ray Evans, Colin Pardoe, Patrick Purcell, Tom Quirk, and David Stockwell, but remain responsible for all remaining errors.

References

- Allison, I., M. Bird, J. Church, M. England, I. Enting, D. Karoly, M. Raupach, J. Palutikof, S. Sherwood 2010. *The Science of Climate Change. Questions and Answers*. Australian Academy of Sciences, Canberra.
- Angrist, J.D. and J-S Pischke (2009). *Mostly harmless econometrics: an empiricist's companion*. Princeton UP, Princeton and Oxford.
- Arrhenius, S. 1896. On the influence of carbonic acid in the air on the temperature on the ground. *Philosophical Magazine and Journal of Science*, Series 5, Volume 41, 237-276.
- Chambers, J., A. Miller, R. Morgan, B. Officer, M. Rayner, T. Quirk 2010. Clearing the air on climate. *Energy and Environment*, 2.16:633-639.

- Curtin, T. 2009a. The Contradictions of the Garnaut Review. *Quadrant*, January-February 2009.
- Curtin, T. 2009b. Climate Change and Food Production. *Energy and Environment*, 20.7
- Dessler, A.E., Z. Zhang, and P. Yang 2008. Water-vapor climate feedback inferred from climate fluctuations 2003-2008. *Geophysical Research Letters*, 34 L20704, 1-4.
- Dessler A.E. and S.M. Davis 2010. Trends in tropospheric humidity from reanalysis systems. *Journal of Geophysical Research*, 115, D19127.
- Dyson, F. 2007. *A Many-colored Glass. Reflections on the Place of Life in the Universe*. University of Virginia Press. Charlottesville and London.
- Farquhar, G.D., S. von Caemmerer, J.A. Berry 1980. A biochemical model of photosynthetic CO₂ assimilation in leaves of C3 species. *Planta*, 149:78-90
- Freedman, David A. and D. Collier, J.S. Sekhon, P.B. Stark (eds.) (2010). *Statistical Models and Causal Inference*. CUP, Cambridge.
- Forster, P. and V. Ramaswamy *et al.* 2007. Changes in Atmospheric Constituents and in Radiative Forcing, in Solomon *et al.* 2007 (chapter 2)
- Gaffen, D.J. and R.J. Ross 1999. Climatology and Trends of U.S. Surface Humidity and Temperature, *Journal of Climate*, vol.12.
- Garnaut, R. 2008. *The Garnaut Climate Change Review. Final Report*. CUP, Cambridge and Melbourne.
- Garnaut, R. 2011. The Science of Climate Change. www.garnautreview.org
- Hansen, J. E., M. Sato, P.Kharecha, and K. von Schuckmann 2011. Earth's Energy Imbalance and Implications. NASA-GISS, New York.
- Hegerl, G. and F. Zwiers *et al.* 2007. Understanding and Attributing Climate Change, in Solomon *et al.* 2007 (chapter 9).
- Kaufmann, R.K. and D.I Stern (2002). Cointegration analysis of hemispheric temperature relations. *J. Geophys. Res.* **107**.
- Kaufmann, R. K., H. Kauppi, and J.H. Stock (2006a). 'Emissions, concentrations, and temperature: A time series analysis', *Climatic Change*, **77**, 3-4.

- Kaufmann, R. K., H. Kauppi, and J. H. Stock (2006b). The relationship between radiative forcing and temperature: what do statistical analyses of instrumental temperature record measure? *Climatic Change* 77: 279–289.
- Kelly, A. 2010. A materials scientist ponders global warming/cooling. *Energy and Environment*, 21.7, October, 611- 632.
- Kiehl, J. T. and K.E. Trenberth 1997. Earth's annual global mean energy budget. *Bulletin of the American Meteorological Society*, 78.2:197-208.
- Kininmonth, W. 2010. Clausius-Clapeyron and the regulation of global warming. *Fisica E*, 26, 5-6, 61-70
- Klotzbach, P. J., R. A. Pielke Sr., R. A. Pielke Jr., J. R. Christy, and R. T. McNider 2009, an alternative explanation for differential temperature trends at the surface and in the lower troposphere, *Journal of Geophysical Research*, 114, D21102, doi:10.1029/2009JD011841.
- Knorr, W. 2009. Is the airborne fraction of anthropogenic CO₂ emissions increasing? *Geophysical Research Letters*, Vol. 36, L21710, doi:10.1029/2009GL040613, 2009.
- Le Quéré, C., J. Canadell, M. Raupach *et al.* 2009. www.globalcarbonproject.org.
- Lim, Wee Ho and M.L. Roderick 2009. *An Atlas of the Global Water Cycle Based on the IPCC AR4 Climate Models*. ANU E-Press, Canberra.
- Long, S.P. 1991. Modification of the response of photosynthetic productivity to rising temperature by atmospheric CO₂ concentrations – has its importance been underestimated? *Plant, Cell & Environment* 14: 729–739.
- Minschwaner K. and A. Dessler 2004. Water Vapor Feedback in the Tropical Upper Troposphere: Model Results and Observations. *Journal of Climate*, 27:1272-1283
- Myhre, G, E.J. Highwood, K.P. Shine, F. Stordal (1998). New estimates of radiative forcing due to well mixed greenhouse gases. *Geophysical Research Letters*, 25.4, 2715-2718
- Norby, R.J. and Y. Luo 2004. Evaluating ecosystem responses to rising atmospheric CO₂ and global warming in a multi-factor world. *New Phytologist* 162: 281–293.
- Nordell, B. and B. Gervet 2009. Global energy accumulation and net heat emission. *International Journal of Global Warming*. 1.1-3:378-391.
- Paltridge, G., A. Arking, and M. Pook (2009), Trends in middle-and upper-level tropospheric humidity from NCEP reanalysis data. *Theoretical.Applied Climatology*, 98, 351–359.

- Pielke, R. snr., *et al.* 2007. Unresolved issues with the assessment of multidecadal global land surface temperature trends. *Journal of Geophysical Research*, 112: 1-26
- Pierrehumbert, R.T., H. Brogniez, and R. Roca 2007. On the Relative Humidity of the Atmosphere, chapter 6 in *The Global Circulation of the Atmosphere*, T. Schneider and A. Sobel, eds., Princeton University Press, 2007.
- Quirk, T. 2011. Carbon dioxide, water vapour and methane (forthcoming).
- Rahmstorf, S., Cazenave A., Church J.A., Hansen J.E., Keeling R.F., Parker D.E., and R.C.J. Somerville. 2007. Recent climate observations compared to projections. *Science*, Vol. 316, No. 709, doi:10.1126/science.1136843.
- Randall, D.A. and R.A. Wood *et al.* 2007. Climate Models and their Evaluation, in Solomon *et al.* 2007 (Chapter 8).
- Raupach, M. R., J. G. Canadell, and C. Le Quéré 2008. Anthropogenic and biophysical contributions to increasing CO₂ growth rate and airborne fraction, *Biogeosciences*, 5, 1601-1613.
- Rocco, F. 2003. *The miraculous fever tree*. HarperCollins, New York.
- Royal Society 2010. *Climate Change. A summary of the science*. Royal Society, London.
- Soares, P.C. 2010, Warming Power of CO₂ and H₂O: Correlations with temperature changes. *International Journal of Geosciences*, 102-112.
- Solomon, S. *et al.* 2007. *Climate Change 2007. The physical science basis*. IPCC, CUP, Cambridge.
- Stern, D.I., and R.K. Kaufmann (1999). Econometric analysis of global climate change. *Environmental Modelling & Software*. 14, 597-605.
- Stern, N. 2007. *The Economics of Climate Change*. CUP, Cambridge.
- Stockwell, D. 2009. Recent climate observations: Disagreement with projections. *Energy and Environment*, 26.4, 595-596.
- Tol, R.S.J. and A.F. de Vos (1998). A Bayesian analysis of the enhanced greenhouse effect. *Climatic Change*, 38, 87-112.
- Tol, R.S.J. and P. Vellinga (1998). Climate change, the enhanced greenhouse effects and the influence of the sun: a statistical analysis. *Theoretical and Applied Climatology*, 61.
- Von Storch, H. and F. Zwiers 1999. *Statistical Analysis in Climate Research*. CUP, Cambridge.

World Nuclear Association (nd). *Is the cooling of power plants a constraint on the future of nuclear power?* WNA, London.